

COLONOSCOPY BOWEL PREPARATION

PLENVU

NAME: _____ **DATE:** _____ **ARRIVAL TIME:** _____

Read these instructions at least **ONE WEEK** before your scheduled appointment so that you understand how to adequately prepare for the colonoscopy procedure. It is very important that you follow all instructions carefully because inadequate bowel preparation can result in missed lesions, increased procedural time, a potential increase in complication rates and may require that your procedure be rescheduled.

In addition, please contact your insurance company to determine if you will have any out-of-pocket cost for your procedure.

If you have any questions about your *procedure preparation* or *medication guidelines*, please contact your scheduler,

_____ at (614) 458-1183, extension # _____.


Procedure Location

Go to Outpatient Registration at the location indicated below.

- | | | |
|---|---|--|
| <input type="checkbox"/> Downtown Endoscopy Center
700 E. Broad St., 1st Floor
Columbus, OH 43215 | <input type="checkbox"/> Westerville Endoscopy Center
300 Polaris Parkway, Suite 1500
Westerville, OH 43082 | <input type="checkbox"/> Riverview Surgery Center
2401 N. Columbus St.
Lancaster, OH 43130 |
| <input type="checkbox"/> Grant Medical Center
Surgical & Heart Entrance
300 E. Town St.
Columbus, OH 43215 | <input type="checkbox"/> Other: _____

_____ | |

Please note: your arrival time is subject to change based upon the facility's schedule. You will receive a reminder call approximately two (2) days before your scheduled procedure at which time your arrival time will be confirmed. Please arrive at the time given to you by the facility.



“A colonoscopy is an exam used to detect changes or abnormalities in the large intestine (colon) and rectum. During a colonoscopy, a long, flexible tube (colonoscope) is inserted into the rectum. A tiny video camera at the tip of the tube allows the doctor to view the inside of the entire colon. If necessary, polyps or other types of abnormal tissue can be removed through the scope...tissue samples (biopsies) can be taken during a colonoscopy as well”.

- MayoClinic.org

Your RX for Plenvu has been sent to Benzer Pharmacy—they complete the prior authorization, if there is one, and apply any discount card available. They will be contacting you on the cost of your prep and will send it directly to your home. If the prescription is too expensive, please inform our office ASAP so that another prep can be sent or a sample can be provided.
Benzer Pharmacy- (810) 230-9900

3-4 days prior to your procedure:


Do not eat corn or beans; their outer layer may take longer for your body to digest.
Do not eat popcorn, seeds, nuts, etc. as they have the potential to damage the colon scope.
Please note: If you eat the above foods, it will **NOT** prevent your procedure from being performed! We just ask that you make every effort to avoid these things.

The day before your procedure:

- **No SOLID FOODS! No RED!**
- Only consume clear liquids such as soup broth, strained fruit juice, coffee, tea, soft drinks, white grape juice, lemonade, etc. You can have jello (no fruit cups or cream) & popsicles (as long as they aren't red!)
- Do not drink red liquids or dark purple (e.g. grape juice); they can stain your colon and hinder visibility
- Avoid dairy products (these take longer to digest)

On the day of your procedure:

- Do not eat or drink anything after midnight the night before, or the day of, your procedure. This includes water.
- Do not take your medications unless instructed to do so (see medication guidelines)



M E D I C A T I O N G U I D E L I N E S

Blood Thinners

Please discuss with your prescribing physician whether or not it is safe for you to discontinue your blood thinners (i.e. Coumadin) for 4 days prior to your procedure. Please inform the Dr and Nurse if you are unable to stop them at the time of your procedure.

Blood Pressure, Seizure, & Heart Medications

These medications may be taken 3 hours prior to your procedure with a small sip of water.

Diabetic Medications

These medications are not to be taken the morning of the procedure. You may bring them with you to take after your procedure or take them when you get home.

Iron Supplements

Iron Supplements must be stopped one (1) week prior to your procedure. This medication makes the colon very dark, so it makes it difficult to see and increases the chances of your physician missing lesions, polyps, etc. Iron infusions do not cause this and are okay to continue.

Non-Steroidal Anti-Inflammatory (NSAID) Medications & Chronic Pain Medications

NSAID medications should be stopped 4 days before your procedure. This includes (but is not limited to) Advil, Aleve, Ascriptin, Celebrex, Empirin, Ibuprofen, Motrin, Naprosyn, Piroxicam, Sulindac, Voltaren.

If you need to take any medication for pain, you may take Tylenol (acetaminophen). If you take chronic pain medications, you may take them the morning of your procedure (3 hours before), but it is preferred that you do not since you will be administered pain medications for your procedure.

If you take any of these medications prior to your procedure, please inform your physician and nurse.

If you are unsure about the type of medication(s) you are taking, please contact your pharmacist.

PLENVU BOWEL PREPARATION INSTRUCTIONS

TAKE DOSE 1 Start Dose 1: *12pm (midnight before proc)*

1. Use the mixing container to mix the contents of Dose 1 pouch with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2-3 minutes. Take your time— slowly finish the dose within 30 minutes.

2. Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes

WAIT APPROXIMATELY 12 HOURS FROM THE START OF DOSE 1, THEN...

TAKE DOSE 2 *4 hour prior to arrival*

1. Use the mixing container to mix the contents of Dose 2 (Pouch A and Pouch B) with at least 16 ounces of water by shaking or using a spoon until it's completely dissolved. This may take up to 2 to 3 minutes. Take your time— slowly finish the dose within 30 minutes.

2. Refill the container with at least 16 ounces of clear liquid. Again, take your time and slowly finish all of it within 30 minutes.

Stay Hydrated!

It's important that you drink clear liquids before, during and after your prep.

*Plenvu can be refrigerated. Use within 6 hours after its mixed with water

BIOPSY RESULTS

If biopsy is taken during your procedure, we will contact you with those results within 7-10 business days after your procedure. If you do not hear from us, please call our office at (614) 458-1183 and dial the corresponding extension for your physician's medical assistant:

Dr. Gordon Kim (ext. 230)
 Dr. Tasos Manokas (ext. 248)
 Dr. Adam Tzagournis (ext. 222)
 Dr. John Tzagournis (ext. 226)
 Dr. Michael Tzagournis (ext. 225)

TRANSPORTATION

For your safety, you **must** have an adult accompany you to the procedure and remain at the facility the entire time. Due to the medications you will be given, you will be unable to work, drive, or take public transportation after your procedure. It is a good idea to arrange for an alternate person to accompany you in the event that your primary support person is unable to do so.